arXiv:1302.0349v1 [math.OA] 2 Feb 2013

QUANTITATIVE K-THEORY AND SPIN CHERN NUMBERS

TERRY A. LORING

Abstract. We validate a method of computing spin Chern numbers that was developed with Hastings and only conjectured to be correct. Specifically, the Pfaffian Bott index can be computed by the "log method" for commutator norms up to a specific constant.

We establish lower bounds on the distance almost commuting unitaries must be moved to change an index. This is done for the Bott index, the winding number index for general unitary matrices, and for the Pfaffian Bott index for self-dual matrices.

1. Introduction

We begin with a survey, and some improved theorems, in the subject of almost commuting unitary matrices and the winding number index discovered with Exel [6]. Our eventual goal is to improve known mathematical results regarding invariants of almost commuting matrices by making them quantitative, so that they be more useful in applications, especially in relation to topological insulators. In the present paper we stick with stability results, with the expectation that in numerical studies of index the averaging over disorder [15, 12, 7] can be done more intelligently. However, we anticipate applications in the classification of C^*-algebras.

The seasoned veterans from the world of C^*-algebras are warned that the Bott index discussed here is the one developed with Hastings to be computable numerically, and it involves different functions in its definition from what is usual [6, Definition 1.2]. That such invariants, contingencies related to stable relations, are not canonically defined is a known difficulty, discussed in [3].

We follow mathematical conventions, so U^* refers to the conjugate-transpose when U is a matrix.

Our main result, from the standpoint of phyics, is that the spin Chern number of a finite system in 2D can be computed as follows. As discussed in length in [12], the position operators, assume the geometry of the model is the torus, are two commuting unitary matrices, but the band-compressed position operators are almost commuting matrices acting on low-energy space that are almost unitary. If the diagonalization of the Hamimtonian is done preserving symmetry, if we start in class AII we end up with matrices A and B so that $A^\sharp = A$ and $B^\sharp = B$ (within machine precision) and only have approximate relation

$$AB \approx BA$$
$$A^*A \approx I$$
$$B^*B \approx I$$

If we compute any approximate self-dual logarithm iK of B, meaning

$$-\pi \leq K \leq \pi$$

2000 *Mathematics Subject Classification.* 19M05, 46L60, 46L80.

Key words and phrases. K-theory, topological insulator, index, operator norm, commutator, C^*-algebra.

and $K^\sharp = K$ and $e^{iK} \approx B$, then the spin Chern number of the system will be even or odd depending on the sign of

$$\mathrm{Pf}\left(Q^* \left(\begin{array}{cc} \frac{1}{\pi}K & \frac{1}{2}\left\{ \sqrt{I - \frac{1}{\pi^2}K^2}, U \right\} \\ \frac{1}{2}\left\{ \sqrt{I - \frac{1}{\pi^2}K^2}, U^* \right\} & -\frac{1}{\pi}K \end{array} \right) Q \right).$$

Where $\{\cdot, \cdot\}$ is the anti-commutator and Q is a specific matrix discussed below that creates anti-symmetry in the formula so that that Pfaffian makes sense.

Of course the logarithm, as a matrix function, will be discontinuous as we cannot assume anything about the spectum of A and B beyond Kramers doubling. Put another way, the various self-dual approximate logarithms of B are far apart, so it is not even clear that the sign of the Pfaffian is well defined. This issue we adrress in the following sections.

There is a separate issue of how to compute approximate logarithms of almost unitary matrices, and how to be sure to get a self-dual output given a self-dual input. That is discussed in a separate paper [14].

In practice, it is feasable to simply compute the unitary parts of A and B before computing further, so

$$U = A\left(A^*A\right)^{-\frac{1}{2}}$$

which is unitar and self-dual, as is

$$V = B\left(B^*B\right)^{-\frac{1}{2}}.$$

The comparison of$\|[U, V]\|$ with $\|[A, B]\|$ is not hard. For the rest of the paper we discuss self-dual matrices. We discuss the Pfaffian-Bott index, which depends only on U and V (or A and B) but was shown [12, Lemma 5.8] to equal the spin Chern index, at least for large systems and other mild assumptions.

Our results on the Pfaffian-Bott index are quantitative, but the correspondance with spin Chern numbers is not quantitative. This will be possible once certaining new quantitiative results concerning localization of Wannier functions or concerning approximation by commuting matrices are developed.

Our results are principally stated in terms of unitary matrices. However, the study of almost commuting unitary element of C^*-algebras is not that different. We know this because we know that the soft-torus is RFD [2]. This needs to be checked for the real soft torus, whose definition is easy enough, now that we have a theory of generator and relations for real C^*-algebras [20].

A unitary U in a C^*-algebra $\mathcal{A}$ might have a logarithm. Given $K^* = K$ in $\mathcal{A}$ so that $U = e^{iK}$ and $\|[U, V]\| \leq \frac{1}{8}$ for a second unitary element V in $\mathcal{A}$, then find that the Bott index in $K_0(\mathcal{A})$ will be determined by

$$\left(\begin{array}{cc} \frac{1}{\pi}K & \frac{1}{2}\left\{ \sqrt{I - \frac{1}{\pi^2}K^2}, U \right\} \\ \frac{1}{2}\left\{ \sqrt{I - \frac{1}{\pi^2}K^2}, U^* \right\} & -\frac{1}{\pi}K \end{array} \right).$$

That is, Theorem 6.2 holds true for all unital C^*-algebras. Indeed, the proof is the same as in the matrix case.

2. Phantom and ordinary almost commuting matrices

Given two unitary matrices U and V with $\delta = \|[U,V]\|$, we find

$$\|VUV^*U^* - I\| = \|[U,V]\|$$

and so by the spectral theorem

$$\sigma\left(VUV^*U^*\right) \subseteq \left\{ z \in \mathbb{T} \,\middle|\, |z-1| \leq \delta \right\}.$$

Thus when $\delta < 2$ we can define $\left(VUV^*U^*\right)^t$ for t between 0 and 1, using a branch of x^t with discontinuity on the negative x-axis. This is a continuous path of unitary matrices from I to VUV^*U^* and

$$\det(VUV^*U^*) = \det(I) = 1$$

so

$$t \mapsto \det\left(\left(UVU^*V^*\right)^t\right)$$

is a loop on the unit circle. We define $\omega(U,V)$ to be the winding number of this path.

This winding number invariant is very computable. There are a few alternate formulas, including

$$\omega\left(U,V\right) = \operatorname{Tr}\left(\frac{1}{2\pi i}\log\left(VUV^*U^*\right)\right) \tag{2.1}$$

which we easily prove: since in some basis VUV^*U^* is diagonal and unitary,

$$VUV^*U^* = \begin{pmatrix} e^{i\theta_1} & & \\ & \ddots & \\ & & e^{i\theta_n} \end{pmatrix}$$

for some $-\pi < \theta_j < \pi$ and

$$\operatorname{Tr}\left(\frac{1}{2\pi i}\log\left(VUV^*U^*\right)\right) = \frac{1}{2\pi i}\operatorname{Tr}\begin{pmatrix} i\theta_1 & & \\ & \ddots & \\ & & i\theta_n \end{pmatrix} = \frac{1}{2\pi}\sum \theta_j$$

and

$$\det\left(\left(VUV^*U^*\right)^t\right) = \det\begin{pmatrix} e^{it\theta_1} & & \\ & \ddots & \\ & & e^{it\theta_n} \end{pmatrix} = \prod e^{it\theta_j}$$

and this path has winding number

$$\omega\left(U,V\right) = \frac{1}{2\pi}\sum \theta_j.$$

Lemma 2.1. *When U and V are commuting unitary matrices, $\omega(U,V) = 0$.*

Proof. In this case the path of determinants is the constant path. ☐

Lemma 2.2. *For*

$$U = \begin{pmatrix} 0 & & & & 1 \\ 1 & 0 & & & \\ & 1 & \ddots & & \\ & & \ddots & 0 & \\ & & & 1 & 0 \end{pmatrix}, \quad V = \begin{pmatrix} e^{i\pi/n} & & & & \\ & e^{2i\pi/n} & & & \\ & & \ddots & & \\ & & & e^{-i\pi/n} & \\ & & & & 1 \end{pmatrix}$$

we have $\omega(U,V)=-1$.

Proof. We find$VUV^*U^*=e^{-\frac{2\pi i}{n}}I$ and so

$$\frac{1}{2\pi i}\mathrm{Tr}\left(\log\left(VUV^*U^*\right)\right)=\frac{1}{2\pi i}\mathrm{Tr}\left(-\frac{2\pi i}{n}I\right)=-1.$$

□

It is easy to modify the example in Lemma 2.2 to get a pair of unitary matrices with $\|[U,V]\|=\delta$ and $\omega(U,V)=n$ for any $0<\delta<2$ and any n.

Theorem 2.3. *Pairs of unitary matrices with* $\|[U,V]\|=\delta$ *and* $\omega(U,V)\neq 0$ *exist for all* $0<\delta<2$. *For such a pair, the distance to a commuting pair of unitary matrices exceeds* 1, *meaning*

$$\|U-U_1\|+\|V-V_1\|>1$$

whenever U_1 *and* V_1 *are unitary matrices with* $U_1V_1=V_1U_1$. *Indeed,*

$$\|U-U_1\|+\|V-V_1\|\geq\sqrt{1-\tfrac{1}{4}\delta^2}.$$

The proof of this will be broken into lemmas and propositions. Theorem 2.3 is a slight improvement on the results obtained with Exel [6].

Lemma 2.4. *Suppose* $\|[U_0,V_0]\|<2$. *If* $\|[U_1,V_1]\|<2$ *and*

$$\omega(U_0,V_0)\neq\omega(U_1,V_1)$$

then for any path U_s,V_s *of unitary matrices, there must be at least one* s_0 *so that*

$$\|[U_{s_0},V_{s_0}]\|=2.$$

Proof. We will use a homotopy argument. If no such s_0 existed then

$$(s,t)\mapsto\det\left(\left(U_sV_sU_s^*V_s^*\right)^t\right)$$

is a homotopy between the path that determines $\omega(U_0,V_0)$ and the path that determines $\omega(U_1,V_1)$. Therefore the winding numbers of these paths are equal. □

Proposition 2.5. *Suppose* $\|[U,V]\|<2$. *If* $\|[U_1,V_1]\|=2$ *or if* $\|[U_1,V_1]\|<2$ *and*

$$\omega(U_1,V_1)\neq\omega(U,V)$$

then

$$\|U-U_1\|+\|V-V_1\|\geq\sqrt{1-\tfrac{1}{4}\|[U,V]\|^2}+\sqrt{1-\tfrac{1}{4}\|[U_1,V_1]\|^2}.$$

Proof. The general case follows from the case $\|[U_1,V_1]\|=2$ by Lemma 2.4. Since this invariant in zero for commuting pairs, this proposition implies Theorem 2.3.

Since -1 is in the spectrum of $V_1U_1V_1^*U_1^*$ then, by a spectral variation result [1], similar to Weyl's estimate on Hermitian operators,

$$\|V_1U_1V_1^*U_1^*-VUV^*U^*\|\geq\mathrm{dist}\left(-1,\sigma\left(VUV^*U^*\right)\right).$$

We also have

$$\|V_1U_1V_1^*U_1^*-VUV^*U^*\|\leq 2\|U-U_1\|+2\|V-V_1\|$$

and

$$\delta=\mathrm{dist}\left(1,\sigma\left(VUV^*U^*\right)\right).$$



Figure 2.1. The solid curve shows the minimum radius in the gap at -1 in with spectrum of VUV^*U^* for unitaries U and V, as a function of $\delta = \|[U,V]\|$. The lower curve show the minimum distance one must go to find a pair with winding number index either undefined or different.

A little basic geometry shows

$$\left(\operatorname{dist}\left(-1,\sigma\left(VUV^*U^*\right)\right)\right)^2 + \left(\operatorname{dist}\left(-1,\sigma\left(VUV^*U^*\right)\right)\right)^2 = 4$$

and so

$$\|U - U_1\| + \|V - V_1\| \geq \sqrt{1 - \tfrac{1}{4}\delta^2}.$$

□

We now get to a difficult question. Is the winding number invariant the only obstruction to closely approximating U and V by commuting unitary matrices? It is important here that we stick with the operator norm in defining "close approximation" as the answers to these sort of questions can change dramatically if considering the Frobenius norm [8, 9, 18, 19]. (In particular, see the discussion in section III in [18].) Results such as this also change dramatically when the matrices come from different symmetry classes, as seen in [16].

There is an answer, but it is only a non-quantitative, nonconstructive result for small δ. This is Theorem 6.15 in [4] and we repeat it here. Also it matters that we are only interested in results for unitaries in $\mathbf{M}_d(\mathbb{C})$ that are *independent of* d [10, 13].

Theorem 2.6. *For any $\epsilon > 0$, there is a δ in $(0,2)$ so that, whenever U and V are unitary matrices in $\mathbf{M}_d(\mathbb{C})$ with $\|[U,V]\| \leq \delta$ and $\omega(U,V) = 0$, there exist unitary matrices U_1 and V_1 in $\mathbf{M}_d(\mathbb{C})$ so that*

$$\|U - U_1\| + \|V - V_1\| \leq \epsilon$$

and $[U_1,V_1] = 0$.

A serious limitation of the invariant $\omega(U,V)$ is that the only known proofs of Theorem 2.6 work with another invariant $\kappa(U,V)$ that is only equal to $\omega(U,V)$ for small δ. This invariant has become known as the Bott index.

Another limitation of $\omega(U,V)$ is that we don't know how to modify it to work in other symmetry classes. For example if we have self-dual unitary matrices, so $U^\sharp = U$ and $V^\sharp = V$, where $\sharp$ is a specific generalized involution detailed below, we find

$$(VUV^*U^*)^\sharp = U^*V^*UV$$

and so generally VUV^*U^* is not self-dual.

3. A DIRECT K-THEORY INVARIANT — THE BOTT INDEX

We need functional calculus of unitary matrices, also called matrix functions in applied mathematics. An example is above where we applied the logarithm to a unitary matrix. Generally speaking, for the functional calculus $f(V)$ to be define for a unitary matrix we need f defined on the circle. One diagonalizes V via another unitary Q and applies f on the diagonal, so

$$V = Q\begin{pmatrix} e^{i\theta_1} & & \\ & \ddots & \\ & & e^{i\theta_d}\end{pmatrix}Q^* \implies f(V) = Q\begin{pmatrix} f\left(e^{i\theta_1}\right) & & \\ & \ddots & \\ & & f\left(e^{i\theta_2}\right)\end{pmatrix}Q^*.$$

However, most of our calculations will involve Fourier series, and traditionally those are defined in terms of scalar functions that are periodic.

Definition 3.1. Assume then that f is periodic of period 2π we define $f[V]$ as $\tilde{f}(V)$ where $\tilde{f} = f \circ \log$. In other words,

$$V = Q\begin{pmatrix} e^{i\theta_1} & & \\ & \ddots & \\ & & e^{i\theta_d}\end{pmatrix}Q^* \implies f[V] = Q\begin{pmatrix} f\left(\theta_1\right) & & \\ & \ddots & \\ & & f\left(\theta_1\right)\end{pmatrix}Q^*.$$

When f has uniformly convergent Fourier series, this is easier:

$$f(x) = \sum_{n=-\infty}^{\infty} a_n e^{inx} \implies f[V] = \sum_{n=-\infty}^{\infty} a_n V^n. \tag{3.1}$$

Definition 3.2. Define

$$f(x) = \frac{1}{128}\left(150\sin(x) + 25\sin(3x) + 3\sin(5x)\right)$$

and g and h:

$$g = \begin{cases} 0 & x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \\ \sqrt{1-f^2} & x \notin \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \end{cases}$$

and

$$h = \begin{cases} \sqrt{1-f^2} & x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \\ 0 & x \notin \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \end{cases}$$

which are shown in Figure 3.1. For any unitaries set

$$B(U,V) = \begin{pmatrix} f[V] & g[V] + \frac{1}{2}\left\{h[V], U\right\} \\ g[V] + \frac{1}{2}\left\{h[V], U^*\right\} & -f[V] \end{pmatrix}$$



FIGURE 3.1. Functions for the standard Bott index (trig method).

and if $\|[U,V]\| \leq 0.206007$ define $\kappa(U,V)$ as the integer

$$\kappa(U,V) = \frac{1}{2}\mathrm{Sig}\left(B(U,V)\right).$$

For an invertible, hermitian matrix A we define its *signature* $\mathrm{Sig}(A)$ as the number (with multiplicity) of positive eigenvalues minus the number of negative eigenvalues. We need to know that $\|[U,V]\| \leq 0.206007$ forces $B(U,V)$ to be invertible.

It should be noted that

$$\begin{aligned}\gamma_1 &= f(\theta_2)\\ \gamma_2 &= g(\theta_2) + h(\theta_2)\cos(\theta_1)\\ \gamma_3 &= h(\theta_2)\sin(\theta_1)\end{aligned}$$

defines the coordinates of a map from $\mathbb{T}^2 \to S^2 \subseteq \mathbb{R}^3$ that has mapping degree one. Also notice $gh = 0$ and $f^2 + g^2 + h^2 = 1$.

Theorem 3.3. *Suppose U and V are unitaries and*

$$\delta = \|[U,V]\| \leq 0.206007.$$

1. *The hermitian matrix* B(U,V) *has a spectral gap at* 0 *of radius at least* $\frac{19}{20}\sqrt{1-5\delta}$*. Indeed, the gap is at least as large as the function of* δ *plotted as a solid curve in Figure 3.2.*
2. *The distance* $\|U - U_1\| + \|V - V_1\|$ *needed so that* $B(U_1,V_1)$ *has* 0 *in its spectrum is at least* $\frac{1}{5}\sqrt{1-5\delta}$*. Indeed, this distance is at least as large as the function of* δ *plotted as a dashed curve in Figure 3.2.*



FIGURE 3.2. In Both: Solid curve on top is the guaranteed spectral gap in $B(U,V)$ for a given δ. The dashed curve is the distance one can move where it is proven the gap will not close. The dotted curve in the left plot is $\frac{19}{20}\sqrt{1-5\delta}$. The dotted curve on the right is $\frac{1}{5}\sqrt{1-5\delta}$.

This will be a lot of work to prove, and the gap is much smaller than we saw for VUV^*U^*. Why do we bother? The point is symmetry.

Suppose $U^\sharp = U$ and $V^\sharp = V$ for unitaries in $\mathbf{M}_{2d}(\mathbb{C})$ and with $\sharp$ the generalized involution discussed in the next section, that physicists call the dual. Then $B(U,V)$ is in $\mathbf{M}_{2d}(\mathbb{C}) \otimes \mathbf{M}_2(\mathbb{C})$ which has on it the generalized involution $\tau = \sharp\otimes\sharp$. In terms of real C^*-algebras, this is a copy of $\mathbf{M}_{2d+2}(\mathbb{C})$ with the transpose operation. In physics language, we are tensoring two half-odd-integer spin systems to get a system with integer spin, in a non-standard basis. We find

$$B(U,V)^\tau = -B(U,V)$$

and so $B(U,V)$ defines a class in

$$K_2(\mathbb{R}) \cong K_{-2}(\mathbb{H}) \cong \mathbb{Z}/2$$

that is computed directly in terms of the a Pfaffian, hence the Pfaffian-Bott index studied in [15]. In return for a small gap, indeed no guaranteed gap if $\|[U,V]\|$ is too large, we get a construction that is amenable to symmetries.

An easy upper bound on the gap radius can be found, using the example in Lemma 2.2, shown in Figure 3.3. This shows we cannot get a big a gap using $B(U,V)$ as was possible with VUV^*U^*, but that the situation is likely not as bad as Figure 3.2 indicates.

Fortunately $B(U,V)$ is readily computable since f, g and h we chosen to have rather fast decay in their Fourier coefficients. Thus we were able to replace (3.1) by the simpler evaluation of order-5 trig polynomials. For applications to index studies, the following is the most useful. We will later have a version of this for the Pfaffian-Bott index.

Proposition 3.4. *Suppose* $\|[U,V]\| \leq 0.206007$*. If* $\|[U_1,V_1]\| \leq 0.206007$ *and*

$$\kappa(U_1,V_1) \neq \kappa(U,V)$$



FIGURE 3.3. Upper bound on the gap radius, as $\|[V,U]\|$ varies from 0 to about 0.21.

then

$$\|U-U_1\|+\|V-V_1\|\geq\frac{19}{20}\sqrt{1-5\,\|[U,V]\|^2}+\frac{19}{20}\sqrt{1-5\,\|[U_1,V_1]\|^2}.$$

Another consequence of Theorem 3.3 is the following.

Theorem 3.5. *Suppose U and V are unitary matrices. If $\|[U,V]\|\leq 0.206007$ then*

$$\kappa(U,V)=\omega(U,V).$$

Proof. Exel [5] showed that for $\|[U,V]\|<2$ the winding invariant equals an abstract K-theory invariant. In the notation of [5], this is defined in terms of b_ϵ in $K_0(A_\epsilon)$. It is easy to check that as long $B(u,v)$ has a spectral gap, where u and v are the generators of the soft torus A_ϵ, the K-theory class of $B(u,v)$ is b_ϵ. □

4. PFAFFIANS

The Pfaffian of skew-symmetric matrices is not the most familiar object, and it it not clear at the outset how it applies to a problem involving self-dual matrices. Let us start by recalling the dual operation.

We fix

$$Z=Z_N=\left[\begin{array}{cc}0 & I\\ -I & 0\end{array}\right]$$

in $\mathbf{M}_{2N}(\mathbb{C})$ and this specifices the dual operation

$$X^\sharp=-ZX^{\mathrm{T}}Z$$

as

$$\left[\begin{array}{cc}A & B\\ C & D\end{array}\right]^\sharp=\left[\begin{array}{cc}D^{\mathrm{T}} & -B^{\mathrm{T}}\\ -C^{\mathrm{T}} & A^{\mathrm{T}}\end{array}\right].$$

When we discuss $\mathbf{M}_2(\mathbf{M}_{2N}(\mathbb{C})) = \mathbf{M}_{2N}(\mathbb{C}) \otimes \mathbf{M}_2(\mathbb{C}) =$ we require the unitary

$$Q = \frac{1}{\sqrt{2}} \left[\begin{array}{cc} I & -iZ \\ iZ & I \end{array} \right] = \frac{1}{\sqrt{2}} \left[\begin{array}{cccc} I & 0 & 0 & -iI \\ 0 & I & iI & 0 \\ 0 & iI & I & 0 \\ -iI & 0 & 0 & I \end{array} \right]$$

which has the convienient property [12, Lemma 1.3]

$$Q^* X^{\sharp\otimes\sharp} Q = \left(Q^* X Q\right)^{\mathrm{T}}.$$

Here

$$\left[\begin{array}{cc} A & B \\ C & D \end{array} \right]^{\sharp\otimes\sharp} = \left[\begin{array}{cc} D^\sharp & -B^\sharp \\ -C^\sharp & A^\sharp \end{array} \right]$$

or

$$\left[\begin{array}{cccc} A_{11} & A_{12} & A_{13} & A_{14} \\ A_{21} & A_{22} & A_{23} & A_{24} \\ A_{31} & A_{32} & A_{33} & A_{34} \\ A_{41} & A_{42} & A_{43} & A_{44} \end{array} \right]^{\sharp\otimes\sharp} = \left[\begin{array}{cccc} A_{44}^{\mathrm{T}} & -A_{34}^{\mathrm{T}} & -A_{24}^{\mathrm{T}} & A_{14}^{\mathrm{T}} \\ -A_{43}^{\mathrm{T}} & A_{33}^{\mathrm{T}} & A_{23}^{\mathrm{T}} & -A_{13}^{\mathrm{T}} \\ -A_{42}^{\mathrm{T}} & A_{32}^{\mathrm{T}} & A_{22}^{\mathrm{T}} & -A_{12}^{\mathrm{T}} \\ A_{41}^{\mathrm{T}} & -A_{31}^{\mathrm{T}} & -A_{21}^{\mathrm{T}} & A_{11}^{\mathrm{T}} \end{array} \right].$$

Recall the Pfaffian is defined for all skew-symmetric, complex $2n$-by-$2n$ matrices by

$$\mathrm{Pf}\left(O \left[\begin{array}{ccccc} 0 & a_1 & & & \\ -a_1 & 0 & a_2 & & \\ & -a_2 & 0 & a_3 & \\ & & -a_3 & 0 & \ddots \\ & & & \ddots & \ddots \end{array} \right] O^{\mathrm{T}} \right) = \det(O) a_1 a_3 \cdots a_{2n-1}.$$

for O real orthogonal. (All skew-symmetric matrices have such a factorization, a modified Hessenberg decomposition.) The essential properites are that

$$\mathrm{Pf}(YXY^{\mathrm{T}}) = \det(Y)\mathrm{Pf}(X)$$

for arbitrary Y, that the Pfaffian varies continuously, and

$$\left(\mathrm{Pf}(X)\right)^2 = \det(X)$$

so is zero exactly on the set of skew-symmetric, singular matrices.

For matrices with the symmetry $X^{\sharp\otimes\sharp} = -X$ we can define a modified Pfaffian

$$\widetilde{\mathrm{Pf}}(X) = \mathrm{Pf}\left(Q^* X Q\right).$$

We still have

$$\left(\widetilde{\mathrm{Pf}}(X)\right)^2 = \det(X)$$

and that this varies continuously. The sign of the Pfaffian can be used to prove a homotopy result, in the same way we use the determinant to detect that the real orthogonal matrices fall into two connected parts.

Proposition 4.1. *Suppose B is in $\mathbf{M}_{4N}(\mathbb{C})$ and $B^* = B$ and $B^{\mathrm{T}} = -B$ and B is invertible. Then*

$$\mathrm{Pf}(B) \in \mathbb{R} \setminus \{0\}.$$

If B_1 and B_2 are elements of

$$\mathcal{H} = \left\{ B \in \mathbf{M}_{4N}(\mathbb{C}) \middle| \, B^* = B = -B^{\mathrm{T}} \textit{ is invertible} \right\},$$

then they can be connected by a path in $\mathcal{H}$ *if and only if with* $\operatorname{Pf}(B_1)$ *and* $\operatorname{Pf}(B_2)$ *have the same sign.*

Proof. We can apply Theorem 8.7 in [12] to iB and learn that there is a real orthogonal matrix O so that $\det(O) = 1$ and

$$B = O \left[\begin{array}{cccccc} 0 & i\lambda_1 & & & & \\ -i\lambda_1 & 0 & & & & \\ & & 0 & i\lambda_2 & & \\ & & -i\lambda_2 & 0 & & \\ & & & & \ddots & \ddots \\ & & & & \ddots & \ddots \end{array}\right] O^{\mathrm{T}}$$

and all the real numbers λ_j are positive except λ_1 which as the same sign as $\operatorname{Pf}(B)$. It is clear from this form that two such matrices with Pfaffian of the same sign will be connected. The spectrum of B is $\{\pm\lambda_1, \ldots, \pm\lambda_{2N}\}$. Thus there will be an even number of negative eigenvalues, so $\det(B) > 0$. Since the square of the Pfaffain is the determinant, we find $\operatorname{Pf}(B)$ is real, and as B is invertible, the Pfaffian cannot be zero. Since the Pfaffian varies continuously, it is not possible to connect two matrices in $\mathcal{H}$ that have Pfaffians of opposite signs. □

Now we explain the Pfaffian-Bott index.

Definition 4.2. Let f, g, h and $B(U,V)$be as in §3. If $\|[U,V]\| \leq 0.206007$ define $\kappa(U,V)$ as the value in $\{\pm 1\}$ given by

$$\kappa_2(U,V) = \operatorname{Sign}\left(\widetilde{\operatorname{Pf}}\left(B(U,V)\right)\right).$$

Lemma 4.3. *When* U *and* V *are commuting unitary matrices,* $\omega(\mathbf{U},\mathbf{V}) = 1$.

Proof. One proof is to use Lemma 6.4 of [11]. Here is another.

One easily checks that κ_2 remains constant along a path so long as $\|[U_t, V_t]\| \leq 0.206007$. One can use functional calculus, and so keep the self-dual condition, to deform a commuting pair U_0 and V_0 that is self-dual over to $U_1 = I$ and $V_1 = I$. One can chen compute that

$$B(I,I) = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}$$

and

$$\widetilde{\operatorname{Pf}}\begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix} = \operatorname{Pf}\begin{bmatrix} iZ & 0 \\ 0 & -iZ \end{bmatrix} = \operatorname{Pf}(iZ)\operatorname{Pf}(-iZ) = i^n(-i)^n = 1.$$

□

Proposition 4.4. *Suppose* $\|[U,V]\| \leq 0.206007$ *and that* U,V, U_1, V_1 *are self-dual unitary matrices. If* $\|[U_1,V_1]\| \leq 0.206007$ *and*

$$\kappa_2(U_1,V_1) \neq \kappa_2(U,V)$$

then

$$\|U - U_1\| + \|V - V_1\| \geq \frac{1}{8}\sqrt{1 - 8\|[U,V]\|^2} + \frac{1}{8}\sqrt{1 - 8\|[U_1,V_1]\|^2}.$$

Proposition 4.5. *Suppose $\|[U,V]\| \leq 0.206007$ and that U,V, U_1, V_1 are self-dual unitary matrices. If U_1 commutes with V_1 and $\kappa_2(U,V) = -1$ then*

$$\|U - U_1\| + \|V - V_1\| \geq \frac{1}{8} + \frac{1}{8}\sqrt{1 - 8\,\|[U,V]\|^2}.$$

We get larger, if more complicated, estimates below.

5. Proofs

We are able to estimate the radius of the spectal gap in $B(U,V)$ by esitmating the norm of $B(U,V)^2 - I$.

Lemma 5.1. *Suppose f, g and h are continuous, real-valued functions that are 2π-periodic, and with*

$$f^2 + g^2 + h^2 = 1$$

and

$$gh = 0.$$

Suppose U and V are unitary matrices and define

$$S = \left[\begin{array}{cc} f[V] & g[V] + \frac{1}{2}\{h[V],U\} \\ g[V] + \frac{1}{2}\{h[V],U^*\} & -f[V] \end{array} \right].$$

Then $S^* = S$ and

$$\left\|S^2 - I\right\| \leq 2\,\|[h[V],U]\| + \|[f[V],U]\|\,.$$

Proof. Since f, g and h are real-valued, the matrices $f[V]$, $g[V]$ and $h[V]$ are hermitian. Let us write f for $f[V]$, etc. We see easily $S^* = S$ and

$$\left[\begin{array}{cc} f & g + \frac{1}{2}\{h,U\} \\ g + \frac{1}{2}\{h,U^*\} & -f \end{array} \right]^2 - \left[\begin{array}{cc} I & 0 \\ 0 & I \end{array} \right] = \left[\begin{array}{cc} A & B \\ B^* & A^* \end{array} \right]$$

where

$$\begin{aligned} A &= f^2 + g^2 - I + \tfrac{1}{4}\{h,U\}\{h,U^*\} + \tfrac{1}{2}g\{h,U^*\} + \tfrac{1}{2}\{h,U\}g \\ &= -h^2 + \tfrac{1}{4}\{h,U\}\{h,U^*\} + \tfrac{1}{2}g\{h,U^*\} + \tfrac{1}{2}\{h,U\}g \end{aligned}$$

and

$$\begin{aligned} B &= fg + \tfrac{1}{2}f\{h,U\} - gf - \tfrac{1}{2}\{h,U\}f \\ &= \tfrac{1}{2}f\{h,U\} - \tfrac{1}{2}\{h,U\}f. \end{aligned}$$

We have

$$\begin{aligned} \left\|S^2 - I\right\| &\leq \left\| \left[\begin{array}{cc} A & 0 \\ 0 & A^* \end{array} \right] \right\| + \left\| \left[\begin{array}{cc} 0 & B \\ B^* & 0 \end{array} \right] \right\| \\ &= \|A\| + \|B\|\,. \end{aligned}$$

Notice $f^2 + g^2 + h^2 = 1$ forces these functions to take value in $[-1,1]$ so $\|f[V]\| \leq 1$, etc. Therefore

$$\begin{aligned} \|A\| &\leq \tfrac{1}{4}\left\|hUhU^* + Uh^2U^* + UhU^*h - 3h^2\right\| + \tfrac{1}{2}\|gU^*h - ghU^* + hUg - Uhg\| \\ &\leq \tfrac{1}{2}\|h\|\,\|Uh - hU\| + \tfrac{1}{4}\left\|Uh^2 - h^2U\right\| + \|g\|\,\|Uh - hU\| \\ &\leq 2\,\|Uh - hU\| \end{aligned}$$

and

$$\begin{aligned}\|B\| &= \tfrac{1}{2}\left\|hfU + fUh - hUf - Ufh\right\| \\ &\leq \tfrac{1}{2}\left\|h\left[f,U\right]\right\| + \tfrac{1}{2}\left\|\left[f,U\right]h\right\| \\ &\leq \left\|\left[f,U\right]\right\|\end{aligned}$$

so

$$\left\|S^2 - I\right\| \leq 2\left\|\left[h,U\right]\right\| + \left\|\left[f,U\right]\right\|.$$

□

Definition 5.2. Suppose f is continuous and 2π-periodic. Following [17] we define $\eta_f : [0,\infty) \to [0,\infty)$ by

$$\eta_f(\delta) = \sup\left\{\left\|\left[f[V],A\right]\right\| \,\middle|\, V \text{ unitary}, \|A\| \leq 1, \|[V,A]\| \leq \delta\right\}$$

where the supremum is taken over all V and A in every unital C^*-algebra.

Once we have a bound on η_f we can use it to bound more than just commutators. Indeed, by [17, Lemma 1.2], for any two unitaries $V_1$and V_2 we have

$$\left\|f[V] - f[V_1]\right\| \leq \eta_f\left(\|V - V_1\|\right).$$

We need a special case of a Lemma in [17].

Lemma 5.3. *Suppose f, g and h are continuous and periodic, and that g is the trigonometric polynomial*

$$g(x) = \sum_{k=-n}^{n} b_k e^{ikx}.$$

then $\eta(\delta) \leq m\delta + b$ where

$$m = \sum_{k=-n}^{n} |kb_k|$$

and

$$b = \max h(x) - \min h(x)$$

We need this only in the case of real-valued functions, in which case b is the diameter of (the range of) h.

A computer algebra package is recommended here, as it shows

$$f(x)^2 + \frac{407}{512}\cos^6(x)\left(1 + \frac{96}{407}\cos(2x) + \frac{9}{407}\cos(4x)\right) = 1$$

which means

$$g(x) = \sqrt{\frac{407}{512}}\cos^3(x)\sqrt{1 + \frac{96}{407}\cos(2x) + \frac{9}{407}\cos(4x)}\left(1 - \chi_{[-\frac{\pi}{2},\frac{\pi}{2}]}(x)\right)$$

$$h(x) = \sqrt{\frac{407}{512}}\cos^3(x)\sqrt{1 + \frac{96}{407}\cos(2x) + \frac{9}{407}\cos(4x)\chi_{[-\frac{\pi}{2},\frac{\pi}{2}]}(x)}$$

A handy formula here is

$$\frac{407}{320}\left(1 + \frac{96}{407}\cos(2x) + \frac{9}{407}\cos(4x)\right) = \left(1 + \frac{15}{40}\cos^2(x) + \frac{9}{40}\cos^4(x)\right)$$

and we get alternate expression for g and h, in particular

$$h(x)=\frac{\sqrt{10}}{4}\cos^3(x)\sqrt{1+\frac{15}{40}\cos^2(x)+\frac{9}{40}\cos^4(x)}\chi_{[-\frac{\pi}{2},\frac{\pi}{2}]}(x).$$

We new bound the derivative of g and h, computing

$$\frac{d}{dx}\left(\frac{\sqrt{10}}{4}\cos^3(x)\sqrt{1+\frac{15}{40}\cos^2(x)+\frac{9}{40}\cos^4(x)}\right)=\frac{p(\sin(x))}{16\sqrt{q(\sin(x))}}$$

where

$$p(x)=30x(x-1)(x+1)(3x^4-10x^2+15)$$

and

$$q(x)=9x^4-33x^2+64.$$

On $[-1,1]$ the max of $p(x)$ is 150 and the min of $q(x)$ is 64 so we find

$$|h'(x)|\leq\frac{150}{128}$$

and the same for g'.

We need h as a cosine or Fourier series so need

$$c_n=\frac{1}{2\pi}\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}}\cos(nx)\sqrt{\frac{407}{512}}\cos^3(x)\sqrt{1+\frac{96}{407}\cos(2x)+\frac{9}{407}\cos(4x)}\,dx.$$

We computed these with numerical integration, and without checking error estimates, in [15]. We compute these a little more carefully here. We find

$$\begin{aligned}c_n&=\frac{1}{2\pi}\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}}\cos(nx)\sqrt{\frac{407}{512}}\cos^3(x)\sum_{k=0}^{\infty}\binom{1/2}{k}\left(\frac{96}{407}\cos(2x)+\frac{9}{407}\cos(4x)\right)^k dx\\&=\sum_{k=0}^{\infty}\frac{1}{2\pi}\sqrt{\frac{407}{512}}\binom{1/2}{k}\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}}\cos(nx)\cos^3(x)\left(\frac{96}{407}\cos(2x)+\frac{9}{407}\cos(4x)\right)^k dx\\&=\sum_{k=0}^{\infty}I_{n,k},\end{aligned}$$

where the $I_{n,k}$ were defined in-line and are easy to compute with a computer algebra package. The convergence here is rather rapid, as

$$\begin{aligned}I_{n,k}&\leq\frac{1}{2\pi}\sqrt{\frac{407}{512}}\binom{1/2}{k}(-1)^k\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}}\left|\cos(nx)\cos^3(x)\left(\frac{96}{407}\cos(2x)+\frac{9}{407}\cos(4x)\right)^k\right|dx\\&\leq\frac{1}{2\pi}\sqrt{\frac{407}{512}}\binom{1/2}{k}\left(-\frac{105}{407}\right)^k.\end{aligned}$$

Letting T_K denote the the Taylor polynomial

$$T_K(x)\approx\sqrt{1+x}$$

n	0	1	2	3	4	5
a_n	0	$-\frac{150i}{256}$	0	$-\frac{25i}{256}$	0	$-\frac{3i}{256}$
b_n	0.202047	-0.179940	0.125655	-0.066010	0.023445	-0.003886
c_n	0.202047	0.179940	0.125655	0.066010	0.023445	0.003886

TABLE 1. These are approximations to the first coefficients in the Fourier expansions of the f, g and h used to define the Bott index. Extend these to negative indices by the rules $a_{-n} = \overline{a_n}$ and $b_{-n} = b_n$ and $c_n = -c_n$.

of degree K expanded at 0, we have

$$\begin{aligned}\sum_{k=K+1}^{\infty} |I_{n,k}| &\leq \frac{1}{2\pi}\sqrt{\frac{407}{512}} \sum_{k=K+1}^{\infty} \binom{1/2}{k} \left(-\frac{105}{407}\right)^k \\ &= \frac{1}{2\pi}\sqrt{\frac{407}{512}} \left(\sum_{k=0}^{\infty} \binom{1/2}{k} \left(-\frac{105}{407}\right)^k - \sum_{k=0}^{K} \binom{1/2}{k} \left(-\frac{105}{407}\right)^k \right) \\ &= \frac{1}{2\pi}\sqrt{\frac{407}{512}} \left(\sqrt{1-\frac{105}{407}} - T_K\left(-\frac{105}{407}\right) \right)\end{aligned}$$

This means we need $K = 7$ to get six digits absolute accuracy:

Using the values in the table to define

$$h_5(x) = \sum_{n=-5}^{5} c_n e^{inx}$$

or

$$h_5(x) = c_0 + 2c_1\cos(x) + 2c_2\cos(2x) + 2c_3\cos(3x) + 2c_4\cos(4x) + 2c_5\cos(5x)$$

and we find,

$$\left| \frac{d}{dx}(h - h_5) \right| \leq \frac{150}{128} + 1.48498 = 2.656855.$$

and so we can estimate to six decimal places the maximum of $|h - h_5|$ by simply plugging in values between $-\pi$ and π with an even spacing of a little less than 10^{-7}. Keeping track of the errors and rounding up, we find

$$\operatorname{diam}(h(x) - h_5(x)) \leq 0.004110$$

and we note

$$\|h_5'\|_F = \sum_{n=-5}^{5} n|c_n| = 1.48498.$$

The other estimates of this sort, for $h_0, \ldots, h_4$, are summarized in Table 2.

We also can use brute force to find

$$\sup_x |h(x) - h_5(x)| \leq 0.002338.$$

Lemma 5.4. *For any unitary matrix V,*

$$\|h_5[V] - h[V]\| \leq 0.002338$$

n	$m =$ bound on $\|h_n'\|_F$	$b =$ bound on diam $(h(x) - h_n(x))$
0	0	1
1	0.359880	0.732237
2	0.862500	0.350141
3	1.258560	0.106619
4	1.446120	0.017509
5	1.48498	0.004110
∞	2.99208	0

TABLE 2. Bounds on η_h as a slope and an offset.

We get the same error estimate on using only b_{-5} through b_5 when numerically computing $g[V]$.

We also use the folklore estimate

$$\|[h[V], U]\| \leq \|h'\|_{\mathrm{F}} \, \|[U, V]\|$$

where

$$\|h\|_{\mathrm{F}} = \sum_{n \in \mathbb{Z}} |nc_n|$$

is in terms of Banach algebra norm $\|\cdot\|_{\mathrm{F}}$ on functions whose Fourier series converge absolutely, where $\|q\|_{\mathrm{F}}$ is the ℓ^1-norm of the Fourier series of q.

Lemma 5.5. For h as in Definition 3.2, we have

$$\|h'\|_{\mathrm{F}} \leq 2.99208.$$

Proof. We check that

$$h'(x) = \frac{-\frac{1}{\sqrt{3256}} \left(45 \cos^4(x) + 60 \cos^2(x) + 120\right)}{\sqrt{1 + \frac{96}{407} \cos(2x) + \frac{9}{407} \cos(4x)}} \sin(x) \cos^2(x) \chi_{[-\frac{\pi}{2}, \frac{\pi}{2}]}(x)$$

and attack this as three factors. It is easy to see

$$\left\| -\frac{1}{\sqrt{3256}} \left(45 \cos^4(x) + 60 \cos^2(x) + 120\right) \right\|_{\mathrm{F}} = \frac{225}{\sqrt{3256}}$$

and the next factor is not so bad, as we see

$$\begin{aligned} \left\| \frac{1}{\sqrt{1 + \frac{96}{407} \cos(2x) + \frac{9}{407} \cos(4x)}} \right\|_F &\leq \sum_{k=0}^{\infty} \left| \binom{-1/2}{k} \right| \left\| \frac{96}{407} \cos(2x) + \frac{9}{407} \cos(4x) \right\|_F^k \\ &= \sum_{k=0}^{\infty} \binom{-1/2}{k} (-1)^k \left(\frac{105}{407} \right)^k \\ &= \frac{1}{\sqrt{1 - \frac{105}{407}}} = \sqrt{\frac{407}{302}}. \end{aligned}$$

We estimate

$$\left\| \sin(x) \cos^2(x) \chi_{[-\frac{\pi}{2}, \frac{\pi}{2}]}(x) \right\|_{\mathrm{F}}$$

n	$m =$ bound on $\|f_n'\|_F$	$b =$ bound on diam $(f(x) - f_n(x))$
0	0	2
1	1.171875	0.21875
2	1.7578125	0.0234375
∞	1.875	0

TABLE 3. Bounds on η_f as a slope and an offset.

as follows. The Fourier series of $-i\sin(x)\cos^2(x)\chi_{[-\frac{\pi}{2},\frac{\pi}{2}]}$ is

$$\ldots\frac{16}{3465\pi},0,\frac{-4}{315\pi},0,\frac{8}{105\pi},\frac{1}{16},\frac{-8}{15\pi},\frac{-1}{16},0,\frac{1}{16},\frac{8}{15\pi},\frac{-1}{16},\frac{-8}{105\pi},0,\frac{4}{315\pi},0,\frac{-16}{3465\pi},\ldots$$

with terms beyond $n = 3$ being given by

$$-\cos\left(\frac{\pi n}{2}\right)\left(\frac{1}{(n-1)^3-4(n-1)}+\frac{1}{(n+1)^3-4(n+1)}\right).$$

Therefore

$$\begin{aligned}
&\left\|\sin(x)\cos^2(x)\chi_{[-\frac{\pi}{2},\frac{\pi}{2}]}\right\|_F\\
&\quad=\frac{1}{4}+\frac{16}{15\pi}+\frac{18}{105\pi}+\frac{4}{\pi}\sum_{n=3}^{\infty}\frac{1}{(2n+1)^3-4\left(2n+1\right)}\\
&\quad\leq\frac{1}{4}+\frac{16}{15\pi}+\frac{18}{105\pi}+\frac{4}{315\pi}+\frac{16}{3465\pi}+\frac{4}{\pi}\int_4^{\infty}\frac{1}{8x^3+12x^2-2x-3}\,dx\\
&\quad=\frac{1}{4}+\frac{16}{15\pi}+\frac{18}{105\pi}+\frac{4}{315\pi}+\frac{16}{3465\pi}+\frac{1}{4\pi}\ln\left(\frac{81}{77}\right)
\end{aligned}$$

and so

$$\|h'\|_{\mathrm{F}}\leq\frac{225}{\sqrt{3256}}\sqrt{\frac{407}{302}}\left(\frac{1}{4}+\frac{16}{15\pi}+\frac{18}{105\pi}+\frac{4}{315\pi}+\frac{16}{3465\pi}+\frac{1}{4\pi}\ln\left(\frac{81}{77}\right)\right)<2.992076$$

□

We approximate f the same way, but this is just arithmetic since f is already a trigonometric polynomial.

Lemma 5.6. *Let f and g and h be as in Definition 3.2. Then*

$$\eta_f(\delta)\leq m\delta+b$$

for each of the values in Table 3, and

$$\eta_g(\delta)\leq m\delta+b$$

and

$$\eta_h(\delta)\leq m\delta+b$$

for each of the values in Table 2.

Let

$$\beta(\delta)=2\eta_h(\delta)+\eta_f(\delta)$$

which is shown in Figure 5.1.



FIGURE 5.1. The function $\beta(\delta)$ that bounds $\|B(U,V)^2 - I\|$ in terms of $\delta = \|[U,V]\|$.

Theorem 5.7. *Suppose U and V are unitary matrices. Then*

$$\left\|B(U,V)^2 - I\right\| \leq \beta\left(\|[U,V]\|\right)$$

and for $\|[U,V]\| \leq 0.206007$ the gap at 0 in the spectrum of $B(U,V)$ has radius at least

$$\sqrt{1-\beta\left(\|[U,V]\|\right)}.$$

The other key thing we have is how $B(U,V)$ varies.

Theorem 5.8. *If U_j and V_j are unitary matrices then*

$$\left\|B\left(U_0,V_0\right) - B\left(U_0,V_0\right)\right\| \leq \beta\left(\|V_0 - V_1\|\right) + \|U_0 - U_1\|$$

and so

$$\left\|B\left(U_0,V_0\right) - B\left(U_0,V_0\right)\right\| \leq \beta\left(\|V_0 - V_1\| + \|U_0 - U_1\|\right)$$

All our main theorems now follow.

6. THE LOG METHOD

An alternate way to compute the Bott index was considered in [6]. One replaces $B(U,V)$ with

$$B_{\mathrm{L}}(U,V) = \begin{pmatrix} \frac{1}{\pi}K & \frac{1}{2}\left\{\sqrt{I - \frac{1}{\pi^2}K^2}, U\right\} \\ \frac{1}{2}\left\{\sqrt{I - \frac{1}{\pi^2}K^2}, U^*\right\} & -\frac{1}{\pi}K \end{pmatrix}$$

where iK is the logarithm of V, meaning $-\pi \leq K < \pi$ and $e^{iK} = V$. Numerical evidence in [15] suggests that, for small commutators, the Pfaffian-Bott index can be computed using $B_{\mathrm{L}}(U,V)$. We validate this here.

Since the logarithm is not continuous, numerical errors will mean we might accidentally compute the wrong branch of logarithm on V, or indeed any logarithm at all.

We note that when q is periodic,

$$q(K) = q[V].$$

Lemma 6.1. *Suppose f, g and h are real-valued Borel functions on $[-\pi,\pi]$ satisfying*

$$f^2 + g^2 + h^2 = 1$$

and

$$gh = 0.$$

Let $q(x) = f(x)h(x)$ and assume further that q and h are continuous and 2π-periodic. Suppose U and V are unitary matrices and $-iK$ is a logarithm of V and define

$$S = \left[\begin{array}{cc} f(K) & g(K) + \frac{1}{2}\{h(K),U\} \\ g(K) + \frac{1}{2}\{h(K),U^*\} & -f(K) \end{array}\right].$$

Then $S^* = S$ and

$$\left\|S^2 - I\right\| \leq (\|g\| + 1)\,\|[h[V],U]\| + \frac{1}{4}\|[h[V],U]\|^2 + \frac{1}{2}\left\|\left[h^2[V],U\right]\right\| + \|[q[V],U]\|.$$

Proof. We write f for $f(K)$ and compute a bit more carefully than before. We find

$$\begin{aligned}
&\frac{1}{4}\left\|hUhU^* + Uh^2U^* + UhU^*h - 3h^2\right\| \\
&\quad = \frac{1}{4}\left\|hUhU^* - h^2 + UhU^*h - Uh^2U^* + 2Uh^2U^* - 2h^2\right\| \\
&\quad = \frac{1}{4}\left\|-[h,U][h,U]^* + 2\left(Uh^2U^* - h^2\right)\right\| \\
&\quad \leq \frac{1}{4}\|[h,U]\|^2 + \frac{1}{2}\left\|\left[h^2,U\right]\right\|
\end{aligned}$$

and

$$\begin{aligned}
\frac{1}{2}\|g\{h,U^*\} + \{h,U\}g\| &= \frac{1}{2}\|gU^*h + hUg\| \\
&= \frac{1}{2}\|g[U^*,h] + [h,U]g\| \\
&\leq \|g\|\,\|[h,U]\|
\end{aligned}$$

and

$$\begin{aligned}
\tfrac{1}{2}\|f\{h,U\} - \{h,U\}f\| &= \tfrac{1}{2}\|fhU - Ufh + fUh - hUf\| \\
&= \tfrac{1}{2}\|2fhU - 2Ufh + fUh - fhU - hUf + Uhf\| \\
&\leq \|fhU - Uhf\| + \tfrac{1}{2}\|fUh - fhU\| + \tfrac{1}{2}\|hUf - Uhf\| \\
&\leq \|fhU - Uhf\| + \tfrac{1}{2}\|Uh - hU\| + \tfrac{1}{2}\|hU - Uh\| \\
&= \|[fh,U]\| + \|[h,U]\| \\
&= \|[q,U]\| + \|[h,U]\|.
\end{aligned}$$

□



FIGURE 6.1. First part of the path, showing f_t then h_t then g_t.



FIGURE 6.2. Second part of the path, showing only f_t then h_t. Here g_t is zero.

Theorem 6.2. *Suppose U and V are unitary matrices, if $\|[U,V]\| \leq \frac{1}{8}$ then for any choice of K with $-\pi \leq K \leq \pi$ and $e^{iK} = V$, there is a path B_t of invertible self-adjoint matrices between $B(U,V)$ and*

$$\begin{pmatrix} \frac{1}{\pi}K & \frac{1}{2}\left\{\sqrt{I - \frac{1}{\pi^2}K^2}, U\right\} \\ \frac{1}{2}\left\{\sqrt{I - \frac{1}{\pi^2}K^2}, U^*\right\} & -\frac{1}{\pi}K \end{pmatrix}$$

and, if U and V are self-dual, then the path may be chosen with the symmetry $B_t^{\sharp\otimes\sharp} = B_t$.

Proof. We can select paths f_t, g_t and h_t from

$$(f_0, g_0, h_0) = (f, g, h),$$

the standard triple, as in Definition 3.2, to (f_1, g_1, h_1) where

$$f_1(x) = \frac{1}{\pi}x,\ g_1(x) = 0,\ h_1(x) = \sqrt{1 - \frac{1}{\pi^2}x^2} \tag{6.1}$$

and the conditions $f^2 + g^2 + h^2 = 1$ and $gh = 0$ hold along the way. This gives us paths of matrices

$$B_t(U,V) = \left[\begin{array}{cc} f_t(K) & g_t(K) + \frac{1}{2}\{h_t(K), U\} \\ g_t(K) + \frac{1}{2}\{h_t(K), U^*\} & -f_T(K) \end{array}\right]$$

with the needed symetries. It remains to show these are invertible. One needs to compute

$$\left(\|g_t\| + 1\right)\eta_{h_t}(\delta) + \frac{1}{4}\left(\eta_{h_t}(\delta)\right)^2 + \frac{1}{2}\eta_{h_t^2}(\delta) + \eta_{q_t}(\delta) \tag{6.2}$$

and check that this takes value less than 1 at $\delta = \frac{1}{8}$. This is a bit much to do by hand, so we instead show by computer that (6.2) takes value less than 0.95 at $\delta = \frac{1}{8}$ for all t in a mesh $t_1, \ldots, t_w$ selected so that

$$\left\|f_{t_j} - f_{t_{j+1}}\right\|_\infty + \left\|g_{t_j} - g_{t_{j+1}}\right\|_\infty + \left\|h_{t_j} - h_{t_{j+1}}\right\|_\infty \leq \sqrt{1 - 0.95} \approx 0.2236.$$

We can keep f_t fixed at

$$f_t(x) = \begin{cases} -1 & -\pi \leq x \leq -\frac{\pi}{2} \\ \frac{1}{128}\left(150\sin(x) + 25\sin(3x) + 3\sin(5x)\right) & -\frac{\pi}{2} \leq x \leq \frac{\pi}{2} \\ 1 & 1\frac{\pi}{2} \leq x \leq \pi \end{cases}$$

while altering g_t from the standard g to 0. The more interesting part of the path interpolates f_t from the above to $\frac{1}{\pi}x$ while keeping $g_t = 0$ and

$$h_t(x) = \sqrt{1 - f_t(x)}.$$

The graphs of the computed bounds are shown in Figure 6.3. These bounds have been rounded up to accomodate the various errors in computing offset terms when applying Lemma 5.3. The errors in computing Fourier coefficients lead to sub-optimal results, but do not need to be accounted for as it is the computed coefficients that are used when applying Lemma 5.3. The analysis of the error bounds is dull and omitted.

It is apparant that that the limitation on the constant in this result comes from the functions used in the log method (6.1). The computed bounds are shown in Figure 6.4. ☐

The computations of the Pfaffian-Bott index are mostly validated by Theorem 6.2. What remains is validaing the algorithms used to compute a self-dual logarithm of a self-dual unitary. This is discussed in [14].

7. ACKNOWLEDGMENTS

The author wishes to thank Matt Hastings, Robert Israel and Fredy Vides for discussions, useful and entertaining. This work was partially supported by a grant from the Simons Foundation (208723 to Loring).



Figure 6.3. Bounds on $\|B_t(U,V)^2 - I\|$.



Figure 6.4. Bounds on $\|B_1(U,V)^2 - I\|$.

References

[1] R. Bhatia and C. Davis, *A bound for the spectral variation of a unitary operator*, Linear and Multilinear Algebra, 15 (1984), pp. 71–76.

[2] S. Eilers and R. Exel, *Finite-dimensional representations of the soft torus*, Proc. Amer. Math. Soc., 130 (2002), pp. 727–731.

[3] S. Eilers and T. A. Loring, *Computing contingencies for stable relations*, Internat. J. Math., 10 (1999), pp. 301–326.

[4] S. Eilers, T. A. Loring, and G. K. Pedersen, *Morphisms of extensions of C^*-algebras: pushing forward the Busby invariant*, Adv. Math., 147 (1999), pp. 74–109.

[5] R. Exel, *The soft torus and applications to almost commuting matrices*, Pacific J. Math., 160 (1993), pp. 207–217.

[6] R. Exel and T. A. Loring, *Invariants of almost commuting unitaries*, J. Funct. Anal., 95 (1991), pp. 364–376.

[7] I. Fulga, F. Hassler, and A. Akhmerov, *Scattering theory of topological insulators and superconductors*, Physical Review B, 85 (2012), p. 165409.

[8] L. Glebsky, *Almost commuting matrices with respect to normalized hilbert-schmidt norm*, (2010).

[9] F. Gygi, J. Fattebert, and E. Schwegler, *Computation of maximally localized Wannier functions using a simultaneous diagonalization algorithm*, Computer Physics Communications, 155 (2003), pp. 1–6.

[10] P. R. Halmos, *Some unsolved problems of unknown depth about operators on Hilbert space*, Proc. Roy. Soc. Edinburgh Sect. A, 76 (1976/77), pp. 67–76.

[11] M. B. Hastings and T. A. Loring, *Almost commuting matrices, localized Wannier functions, and the quantum Hall effect*, J. Math. Phys., 51 (2010), p. 015214.

[12] M. B. Hastings and T. A. Loring, *Topological insulators and C^*-algebras: Theory and numerical practice*, Ann. Physics, 326 (2011), pp. 1699–1759.

[13] H. Lin, *Almost commuting selfadjoint matrices and applications*, in Operator algebras and their applications (Waterloo, ON, 1994/1995), vol. 13 of Fields Inst. Commun., Amer. Math. Soc., Providence, RI, 1997, pp. 193–233.

[14] T. A. Loring, *Computing a logarithm of a unitary matrix with general spectrum.* arXiv:1203.6151.

[15] T. A. Loring and M. B. Hastings, *Disordered topological insulators via C^*-algebras*, Europhys. Lett. EPL, 92 (2010), p. 67004.

[16] T. A. Loring and A. P. W. Sørensen, *Almost commuting unitary matrices related to time reversal.* arXiv:1107.4187, 2011.

[17] T. A. Loring and F. Vides, *Estimating norms of commutators.* arXiv:1301.4252.

[18] N. Marzari, I. Souza, and D. Vanderbilt, *An introduction to maximally-localized Wannier functions*, Highlight the Month, Psi-K Newsletter, 57 (2003), pp. 129–68.

[19] A. Ruhe, *Closest normal matrix finally found!*, BIT, 27 (1987), pp. 585–598.

[20] A. Sørensen, *Semiprojectivity and the geometry of graphs*, PhD thesis, University of Copenhagen, 2012. www.math.ku.dk/noter/filer.

Department of Mathematics and Statistics, University of New Mexico, Albuquerque, NM 87131, USA.